



PSICOLOGÍA DEL DESARROLLO: INFANCIA Y ADOLESCENCIA

## BEHAVIORAL AND COGNITIVE FUNCTIONING AFTER TRAUMATIC BRAIN INJURY

**Francesca Tebaldi\*, Anna Cantagallo°**

Modulo di Neuropsicologia Riabilitativa – UOMR -

Dipartimento di Neuroscienze/Riabilitazione, Azienda Ospedaliero-Universitaria di Ferrara

\*Psychologist, PhD

° Psychiatrist, Neurologist and Neuropsychologist

### ABSTRACT:

*Background:* Behavioral and cognitive disorders often occur after Traumatic Brain Injury (TBI). Controversial results about the relation between these outcomes are reported.

*Aims:* We aimed: explore: 1) to explore the relation between different aspects of behavioral functioning of TBI patients; 2) to explore the cognitive functioning of patients with behavioral disorders compared to patients without behavioral disorders; 3) to correlate the behavioral and cognitive functioning for both groups.

*Methods:* A group of 74 severe TBI patients admitted in the Neuropsychological Rehabilitation Unit of our Hospital took part in our study [inclusion criteria: LCF  $\geq 6$ ; time post-injury  $\leq 9$  years; no history of neurological or psychiatric disorders]. We assessed the patients' cognitive functioning with tests commonly used in clinical neuropsychology and the behavioral functioning with the Functional Assessment Measure.

*Results:* 1) We found significant correlation between "emotional controls", "interpersonal relationships" and "awareness"; 2) patients with behavioral disorders compared to patients without behavioral disorders show a worse performance in tests for attention and executive functions and a different relation between cognitive and behavioral functioning; 3) the behavioral functioning significantly correlates with attention and executive function only for patients without behavioral disorders, underlying a diffuse disorganization of these cognitive processes for patients with behavioral disorders.

**KEYWORDS:** behavioral functioning, neuropsychological functioning, traumatic brain-injury

### INTRODUCTION

Traumatic Brain Injury (TBI) often leads to cognitive and behavioral disorders and it is a leading cause of disability. During the acute phase, behavioral disorders may be determined by organic factors such as severity of the trauma, but later in post-acute phase they can persist and be aggravated by environmental and social factors, too (Franulic et al., 2000).

Research has suggested that emotional and behavioral difficulties are the most common complaints



## BEHAVIORAL AND COGNITIVE FUNCTIONING AFTER TRAUMATIC BRAIN INJURY

5 years post-injury, regardless of patients' age (Skell et al., 2000; Deb and Burns, 2007) and may be the most critical factor determining the quality of long-term recovery (Warrier et al., 2003).

Several studies have explored the association between different aspects of behavioral functioning in TBI patients. However, it must be noted that they differ in sample size, severity of trauma, time interval between trauma and assessment, assessment and disorder classification procedures, so they cannot be compared.

Two different results about the relation between self-awareness and emotional control have been recently reported:

1. Sawhchyn et al. (2005) administered to patients and relatives the Patient Competency Rating Scale (PCRS) (Prigatano et al., 1986), and the Katz Adjustment Scale-Revised (Jackson et al., 1992). Both scales refer to self-awareness and cognitive, behavioral and emotional functioning. They found less awareness of deficit in patients with emotional adjustment difficulties.
2. Malec et al. (2007) found a direct relation between self-awareness and depression measured with the Neurobehavioral Functioning Inventory (NFI) Depression Scale (Kreutzer et al., 1999). They conceptualized that impaired self-awareness may serve as a barrier to the development of depression.

A different research issue concerns the association between behavioral and cognitive functioning. Some recent studies reported that behavioral functioning is related with executive functions and attention:

1. Hart et al. (2005) combined eight clinical measures of executive functions in a composite score, the Executive Composite (EC). They found that a worse EC score was associated with impaired self-awareness of behavior and attention deficits, estimated using self and significant other questionnaires.
2. Krpan et al. (2007) reported that a lower executive performance was related to the use of "emotion focused coping" in interpersonal relationships, considered more maladaptive compared to "problem focused coping".
3. Baddeley et al. (1997) shared out patients without behavioral disorders from patients with dysexecutive behavioral disorders, estimated with questionnaires administered to patients and caregivers. They found a lower ability of divided attention in patients with behavioral disorders.

To investigate the behavioral and cognitive functioning after TBI, we shared out patients in two groups, depending on their behavioral functioning level:

- "Low functioning" group (if at least one of the behavioral F.A.M. item was  $\leq 5$ )
- "High functioning" group (if every behavioral F.A.M. item was 6 or 7).

### AIMS AND HYPOTHESIS:

- **Aim 1:** to further explore the relation between different aspects of behavioral functioning for the two groups.

**Hypothesis 1:** a direct correlation between interpersonal relationships, emotional control and awareness (behavioral F.A.M. items) for both groups is expected.

- **Aim 2:** to explore the differences in demographic (age, educational level) and clinical data (GCS, coma duration, PTA duration, time post-injury, total F.A.M.) between the two groups.

**Hypothesis 2:** no difference between groups is expected.

- **Aim 3:** to explore the differences in cognitive functioning between the two groups. **Hypothesis 3:** a lower performance in attention and executive function abilities is expected for patients with behavioral disorders.

- **Aim 4:** To correlate the behavioral and cognitive functioning for the two groups.

**Hypothesis 4:** for both groups, a direct correlation of the behavioral F.A.M. items with attention and executive functions abilities is expected.



## PSICOLOGÍA DEL DESARROLLO: INFANCIA Y ADOLESCENCIA

## METHODS

## SUBJECTS

We identified 74 TBI patients (58 males, 16 females) admitted to our Hospital between 2000 and 2008. All patients had a severe TBI in accordance to Glasgow Coma Scale (GCS), coma duration and Post-Traumatic Amnesia (PTA) duration. Inclusion criteria were:

- not younger than 15 and not older than 80 years;
  - Level of Cognitive Functioning  $\geq 6$ ;
  - no linguistic disorders;
  - no motor, visual and auditory disorders which could prevent the neuropsychological tests' administration;
  - no history of previous neurological disease;
  - no history of alcoholism, drug assumption, psychiatric disorders and mental retardation.
- Demographic and clinical data of the two groups are shown in Table I.

|                  | Group   | M      | SD    | Range     | Sig. (p =) |
|------------------|---------|--------|-------|-----------|------------|
| Age              | Low F.  | 32.06  | 14.53 | 15- 73    | .806       |
|                  | High F. | 31.16  | 15.48 | 16- 72    |            |
| Ed. level        | Low F.  | 11.24  | 3.10  | 5 - 18    | .265       |
|                  | High F. | 12.16  | 3.70  | 8 - 20    |            |
| GCS              | Low F.  | 6.05   | 3.17  | 3 - 14    | .907       |
|                  | High F. | 5.94   | 2.35  | 3 - 13    |            |
| Coma duration    | Low F.  | 30.33  | 30.34 | 1 - 120   | .171       |
|                  | High F. | 16.18  | 15.61 | 1 - 45    |            |
| PTA duration     | Low F.  | 11.96  | 7.50  | 1 - 28    | .727       |
|                  | High F. | 11.06  | 8.58  | 1 - 30    |            |
| F.A.M.           | Low F.  | 177.08 | 30.83 | 75 - 210  | .484       |
|                  | High F. | 182.24 | 27.64 | 115 - 210 |            |
| Time post-injury | Low F.  | 13.96  | 23.68 | 1 - 180   | .177       |
|                  | High F. | 7.36   | 6.42  | 2 - 28    |            |

**TAB. I. Demographic and clinical data of the two groups.**

## ASSESSMENT OF BEHAVIORAL FUNCTIONING

F.A.M. is a 30 items, seven levels ordinal scale used in the rehabilitation of any neurological, musculoskeletal and other disorders to assess areas of dysfunction in everyday activities. It is administered by the rehabilitation team in collaboration with the patients and his family. Items are classified in motor functions (16 items related to self-care, sphincter control, mobility, locomotion) and cognitive functions (14 items related to communication, psychosocial adjustment, cognitive functioning) (Hall et al., 2001; Tesio e Cantagallo, 1998).

The general coding is:

- 7 = Complete independence (timely, safely);
- 6 = Modified independence (extra time, device);
- 5 = Supervision;
- 4 = Minimal assist (independence in >75% of task);
- 3 = Moderate assist (independence in 50-75% of task);



## BEHAVIORAL AND COGNITIVE FUNCTIONING AFTER TRAUMATIC BRAIN INJURY

2 = Maximal assist (independence in 25-50% of task);

1 = Total assist (independence in < 25% of task).

We assessed the patient's behavioral functioning with 3 cognitive F.A.M. items concerning behavior: "emotional control", "interpersonal relationships", and "awareness". They respectively refers to:

- **Emotional control:** frequency and severity of depression, anxiety, frustration, apathy, agitation episodes; effect of these episodes on everyday activities; patient's ability to cope with his emotional behavior.
- **Interpersonal relationships:** ability to interact with other people in clinical and social context and to integrate individual needs with other people needs; inappropriate behaviors refer to yell out, to use a scurrilous language, to react aggressively, to show excessive laugh or crying and to show changeable attitudes or inability to get in touch with others.
- **Awareness:** awareness and acceptance of disability, motivation to learn compensatory strategies; to take appropriate safety measures and to have realistic expectations about the future.

## ASSESSMENT OF COGNITIVE FUNCTIONING

### Attention

- Visual Scanning (Matrices) (Spinnler and Tognoni, 1987), to assess selective attention and visual search ability: execution time and number of errors are recorded.

- Trail Making Test (TMT), subtest A e B (Giovagnoli et al., 1996), to assess the cognitive flexibility implied in the execution of two alternating tasks: time execution in part A and B and the difference between the two have been analyzed.

- Verbal span (disyllabic words) (Spinnler and Tognoni, 1987), to assess verbal short-term memory;

- Spatial Span (Corsi) (Spinnler and Tognoni, 1987), to assess visual-spatial short-term memory;

- Immediate Visual Memory (IVM) (Caltagirone et al, 1995).

### Executive Functions

- Wisconsin Card Sorting Test (WCST) (Heaton et al., 2000), to assess the ability to formulate abstract concepts and to flexibly modify strategies depending on environmental modifications. Global Score, Perseverative Response, Non-Perseverative Errors and Failure to maintain a set have been analyzed;

- Behavioral Assessment of Dysexecutive Syndrome (BADS) (Wilson et al., 1996), to assess the ability of actions and routes planning, temporal judgments and cognitive flexibility; we analyzed each subtest's score (Rule shift card, Action program, Key search, Temporal judgment, Zoo map, Modified six elements);

- Raven Progressive Matrices (38) (Spinnler and Tognoni, 1987), to assess the abstract reasoning ability.

### Memory and Learning

- Verbal selective reminding test (Buschke-Fuld) (Spinnler and Tognoni, 1987), to explore whether learned verbal stimuli are stored in short-term or long-term memory (coding and recovery LT processes). We analyzed the following scores: Total recall, Consistent recall and Delayed recall;

- Spatial Sovra-Span (Spinnler and Tognoni, 1987), to assess visual-spatial learning;

- Prose memory (Capitani et al., 1994), to assess both the ability to recall the text regardless of the items' significance (No hierarchic score) and the ability to recall the more important events (Hierarchic score).



## PSICOLOGÍA DEL DESARROLLO: INFANCIA Y ADOLESCENCIA

## STATISTICAL ANALYSIS

An oneway ANOVA was carried out to explore if the two groups were different in relation to:

- Demographic data: age, educational level;
- Clinical data of both the acute phase (GCS, coma duration, PTA duration) and post-acute phase (time post-injury, total F.A.M.);
- Performance in neuropsychological tests.

For the two groups, Spearman's rank correlation was computed:

- Between behavioral F.A.M. behavioral items;
- Between behavioral functioning (behavioral F.A.M items) and cognitive functioning (performance in neuropsychological tests).

Alpha < 0.05 was taken as significance level.

## RESULTS

**Hypothesis 1 confirmed:** A Significant correlations between interpersonal relationships, emotional control and awareness (behavioral F.A.M). items resulted for both groups, as shown in Table II.

|                   | Interpersonal relationships |         | Emotional control |         |
|-------------------|-----------------------------|---------|-------------------|---------|
|                   | Low F.                      | High F. | Low F.            | High F. |
| Emotional control | .691**                      | .611**  |                   |         |
| Awareness         | .550**                      | .695**  | .554**            | .792**  |

Significant two-tailed correlations are pointed up with asterisks (\* for significant value at the .01 level, \*\* for significant value at the .05 level).

**TAB. II. Correlations between the behavioral F.A.M. items for the two groups.**

**Hypothesis 2 confirmed:** Any significant differences between groups was found in demographic and clinical data (see Tab. I).

**Hypothesis 3 confirmed:** The Low functioning group had a worse performance compared to the High functioning group in neuropsychological tests assessing attention and executive function, as Table III reported. Specifically, they seem to have less ability to select attention on target stimuli, to shifting of attention between two alternating tasks and to plan the execution of a complex action sequence within a specific time interval.

**BEHAVIORAL AND COGNITIVE FUNCTIONING AFTER TRAUMATIC BRAIN INJURY**

|                              | Group   | M      | SD    | Range          | Sig. (p=) |
|------------------------------|---------|--------|-------|----------------|-----------|
| <b>Matrices</b>              | Low F.  | 38.33  | 8.66  | 15.25 - 51.75  | .034      |
|                              | High F. | 42.85  | 7.61  | 29.25 - 56.25  |           |
| <b>TMT A</b>                 | Low F.  | 64.65  | 23.11 | 30 - 143       | .02       |
|                              | High F. | 52.36  | 15.87 | 24 - 85        |           |
| <b>TMT B</b>                 | Low F.  | 168.93 | 54.54 | 50 - 280       | .038      |
|                              | High F. | 139.65 | 50.98 | 42 - 303       |           |
| <b>TMT B-A</b>               | Low F.  | 111.31 | 45.54 | 27 - 227       | .034      |
|                              | High F. | 86.57  | 40.92 | 14 - 220       |           |
| <b>BADS<br/>(6 elements)</b> | Low F.  | 83.39  | 20.12 | 46.6 - 108.87  | .001      |
|                              | High F. | 106.04 | 9.39  | 77.73 - 108.87 |           |

TAB. III. Significant differences between groups in the cognitive functioning.

No difference between groups was found in other tests.

**Hypothesis 4 partially confirmed:** As shown in Table IV, behavioral F.A.M. items correlates with tests assessing attention and executive functions almost only for patients with high behavioral functioning.

|                    | <b>Interpersonal relationships</b> |         | <b>Emotional control</b> |         | <b>Awareness</b> |         |
|--------------------|------------------------------------|---------|--------------------------|---------|------------------|---------|
|                    | Low F.                             | High F. | Low F.                   | High F. | Low F.           | High F. |
| <b>TMT A</b>       | .152                               | -.552** | .227                     | -.603** | .025             | -.589** |
| <b>TMT B</b>       | .129                               | -.499*  | -.034                    | -.588** | -.077            | -.699** |
| <b>TMT B-A</b>     | .048                               | -.479*  | -.25                     | -.540** | -.191            | -.659** |
| <b>BADS (Rule)</b> | .247                               | .787**  | .337                     | .664*   | .444*            | .885**  |
| <b>Raven ('38)</b> | -.086                              | .501*   | -.167                    | .569**  | .039             | .647**  |

Significant two-tailed correlations are pointed up with asterisks (\* for significant value at the .01 level, \*\* for significant value at the .05 level).

TAB. IV. Significant correlations between behavioral and cognitive functioning for the two groups.  
**DISCUSSION AND CONCLUSIONS**

**Aim 1:** The strong correlation between awareness, interpersonal relationships and emotional resulted for both groups, suggest that a single, comprehensive process is responsible for all the F.A.M. items



## PSICOLOGÍA DEL DESARROLLO: INFANCIA Y ADOLESCENCIA

concerning behavioral functioning. Our results is in accordance with Sawhchyn et al. (2005) and Malec et al. (2007) findings.

**Aim 2:** Behavioral disorders in TBI patients seemed not related either to demographic data, clinical data of the acute-phase and clinical data of the post-acute phase. As proposed by Franulic et al. 2000, it would be useful to consider the role held by environmental and social factor in the development of behavioral disorders.

**Aim 3:** In accordance with other author's results (Hart et al., 2005; Krpan et al., 2007; Baddeley et al., 1997), patients with behavioral disorders perform worse compared to patients without behavioral disorders in test for attention and executive functions. Further research is needed to determine whether the behavioral functioning contribute in a causal fashion to attention and executive functions abilities.

**Aim 4:** Behavioral functioning correlates with attention and executive functions almost only for patients with high behavioral functioning, suggesting that behavioral disorders are related with a diffuse disorganization of these cognitive processes. It would be useful to explore if some aspects of emotional control, interpersonal relationships, awareness are more responsible than others for supporting cognitive functioning.

Our findings support that a rehabilitative approach focalizing on both cognitive and behavioral functioning would be useful for TBI patients. Nevertheless, only few studies are reported about this issue. Rath et al. (2003) proposed an innovative group treatment for the problem-solving deficits. It incorporated strategies for addressing underlying emotional self-regulation and logical thinking/reasoning deficits, focalizing on both motivational, attitudinal and affective processes and problem-solving skills in patients with TBI. Results showed an improvement in problem solving as assessed using a variety of measures, including (1) executive function (2) problem-solving self-appraisal, (3) self-appraised emotional self-regulation and clear thinking, (4) objective observer ratings of role-played scenarios. An adaptation of this protocol to the Italian context is in progress, (Cantagallo et al., in progress) expecting it could improve behavioral functioning and executive functions abilities in TBI patients.

## REFERENCES

- Baddeley, A., Della Sala, S., Papagno, C., Spinnler, H. (1997). Testing Central Executive Functioning with a Pencil and Paper Test. *Psychology Press*, 3: 62-69.
- Caltagirone, C., Gainotti, G., Carlesimo, G.A., Parnetti, L., ed il Gruppo per la standardizzazione della Batteria per il Deterioramento Mentale (1995). Batteria per la valutazione del deterioramento mentale. Parte 1: Descrizione di uno strumento di diagnosi neuropsicologica. *Archivio di Psicologia, Neurologia e Psichiatria*, 56: 461-470.
- Cantagallo, A., Vestri, A., Guerra, P., Mondin, E., Tebaldi, F., Diller, L. (in progress). Problem Solving Rehabilitation: the Italian project.
- Capitani E., Della Sala S., Laiacona M., Marchetti C., Spinnler H. (1994). Standardizzazione ed uso clinico di un test di memoria di prosa. *Bollettino di Psicologia Applicata*, 209: 47-63.
- Deb, S., Burns, J. (2007). Neuropsychiatric consequences of traumatic brain injury: a comparison between two age groups. *Brain injury*, 21(3): 301-301.
- Della Sala, S., Baddeley, A., Papagno C., Spinnler, H. (1995). Dual-task paradigm: a means to examine the central executive. *Annals of the New York Academy of Sciences*, 769: 161-71.
- Franulic, A., Horta, E., Maturana, R. et al. (2000). Organic personality disorders after traumatic brain injury: cognitive, anatomic and psychosocial factors. A 6-month follow-up. *Brain Injury*, 5:431-439.
- Giovagnoli, A.R., Del Pesce, M., Mascheroni, S., Simoncelli, M., Laiacona, M., Capitani, E. (1996). Trail Making Test: normative values from 287 normal adult controls. *Italian Journal of Neurological Sciences*, 17: 305-309.
- Hall, K.M., Bushnik, T., Lakisic-Kazazic, B., Wright, J., Cantagallo, A. (2001). Assessing traumatic brain injury outcome measures for long-term follow-up of community-based individuals. *Archives of*

**BEHAVIORAL AND COGNITIVE FUNCTIONING AFTER TRAUMATIC BRAIN INJURY**

- Physical Medicine and Rehabilitation, 82:367-74.
- Hart, T., Whyte, J., Kim, J., Vaccaro, M. (2005). Executive function and self-awareness of "Real word" behavior and attention deficits following traumatic brain injury. *Journal of Head Trauma Rehabilitation*, 20(4): 333-347.
- Heaton, R.K., Chelune, G.J., Talley, J.L., Kay, G.G., Curtiss, G. (2000). WCST: Wisconsin Card Sorting Test. Firenze: Giunti-Organizzazioni Speciali. (Edizione italiana a cura di Hardoy M.C., Carta M.G., Hardoy M.J., Cabras P.L.).
- Jackson, H.F., Hopewell C.A., Glass, C.A., Warburg, R., Dewey, M., Ghadiali, E. (1992). The Katz Adjustment Scale: modification for use with victims of traumatic brain and spinal injury *Brain Injury*, 6(2): 109-127.
- Kreutzer J, Seel R, Marwitz J. (1999). *The Neurobehavioral Functioning Inventory*. San Antonio: The Psychological Corp.
- Malec, J.F., Testa, J.A., Rush, B.K., Brown, A.W., Moessner, A.M. (2007). Self-assessment of impairment, impaired self-awareness, and depression after traumatic brain injury. *Journal of Head Trauma Rehabilitation*, 22(3):156-166.
- Prigatano, G.R., Fordyce, D.J., Zeiner, H.K., Roueche, J.R., Pepping, M., Woods, B. (1986). *Neuropsychological rehabilitation after brain injury*. Baltimore: The Johns Hopkins University Press.
- Rath, J.F., Simon, D., Langenbahn, D.M., Sherr, R.L., Diller, L. (2003). Group treatment of problem-solving deficits in outpatient with traumatic brain injury: a randomized outcome study. *Neuropsychological Rehabilitation*, 13(4). 461-488.
- Skell, R.L., Johnstone, B., Schopp, L. Shaw, J., Petroski, G.F. (2000). Neuropsychological predictors of distress following traumatic brain injury.
- Spinnler, H., Tognoni, G. (1987). Standardizzazione e Taratura Italiana di Test Neuropsicologici. *Italian Journal of Neurological Sciences*, 8: supplementum to 6.
- Tesio, A., Cantagallo, A. (1998). The Functional Measure (F.A.M.) in closed traumatic brain injury outpatients: a Rasch-based validation study." *Journal of Measurement Outcome*, 2:79-96.
- Warrier, E.M., Rourke, B.P., Velikonja, D., Metham, L. (2003). Subtypes of emotional and behavioral sequelae in patients with traumatic brain injury. *Journal of Clinical and Experimental Neuropsychology*, 25(7):904-917.
- Wilson, B., Alderman, N., Burgess, P., et al (1996) *Behavioral Assessment of the Dysexecutive Syndrome BADS*. Thames Valley Test Company.

*Fecha de recepción: 28 febrero 2009*

*Fecha de admisión: 19 marzo 2009*